

Maximal inequalities and a law of the iterated logarithm for negatively associated random fields*

LI-XIN ZHANG

Department of Mathematics, Zhejiang University, Hangzhou 310027, China

Abstract

The exponential inequality of the maximum partial sums is a key to establish the law of the iterated logarithm of negatively associated random variables. In the one-indexed random sequence case, such inequalities for negatively associated random variables are established by Shao (2000) by using his comparison theorem between negatively associated and independent random variables. In the multi-indexed random field case, the comparison theorem fails. The purpose of this paper is to establish the Kolmogorov exponential inequality as well a moment inequality of the maximum partial sums of a negatively associated random field via a different method. By using these inequalities, the sufficient and necessary condition for the law of the iterated logarithm of a negatively associated random field to hold is obtained.

Key Words: negative association, law of the iterated logarithm, random field, Kolmogorov exponential inequality, the maximum partial sums.

Abbreviated Title: LIL for NA Fields

AMS 2000 subject classifications. Primary 60F15

*Research supported by National Natural Science Foundation of China

1 Introduction and the law of iterated logarithm

A finite family of random variables $\{X_i; 1 \leq i \leq n\}$ is said to be negatively associated if for every pair of disjoint subsets A and B of $\{1, 2, \dots, n\}$,

$$\text{Cov}\{f(X_i; i \in A), g(X_j; j \in B)\} \leq 0 \quad (1.1)$$

whenever f and g are coordinatewise non-decreasing and the covariance exists. An infinite family is negatively associated if every finite subfamily is negatively associated. The concept of the negative association was introduced by Alam and Saxena (1981) and Joag-Dev and Proschan (1983). As pointed out and proved by Joag-Dev and Proschan (1983), a number of well-known multivariate distributions possess the NA property. Many investigators discuss the properties and limit theorems of negatively associated random variables. We refer to Joag-Dev and Proschan (1983) for fundamental properties, Newman (1984) for the central limit theorem, Matula (1992) for the three series theorem, Su, et al. (1997) for the moment inequality, Roussas (1996) for the Hoeffding inequality, and Shao (2000) for the Rosenthal-type maximal inequality and the Kolmogorov exponential inequality. Shao and Su (1999) established the law of the iterated logarithm for negatively associated random variables with finite variances.

Theorem A *Let $\{X_i; i \geq 1\}$ be a strictly stationary negatively associated sequence with $EX_1 = 0$, $EX_1^2 < \infty$ and $\sigma^2 := EX_1^2 + 2 \sum_{i=2}^{\infty} E(X_1 X_i) > 0$. Let $S_n = \sum_{i=1}^n X_i$. Then*

$$\limsup_{n \rightarrow \infty} \frac{S_n}{(2n \log \log n)^{1/2}} = \sigma \quad a.s. \quad (1.2)$$

Here and in the sequel of this paper, $\log x = \ln(x \vee e)$

Let $\{X_{\mathbf{n}}; \mathbf{n} \in \mathbb{N}^d\}$ be a field of random variables, where $d \geq 2$ is a positive integer, $\mathbb{N}^d$ denotes the d-dimensional lattice of positive integers. Through this paper, for $\mathbf{n} = (n_1, \dots, n_d) \in \mathbb{N}^d$, $\mathbf{k} = (k_1, \dots, k_d) \in \mathbb{N}^d$ and $m \in \mathbb{N}$, we denote $|\mathbf{n}| = n_1 \cdots n_d$, $\|\mathbf{n}\| = n_1 + \dots + n_d$, $\mathbf{k}\mathbf{n} = (k_1 n_1, \dots, k_d n_d)$ and $\mathbf{k}m = (k_1 m, \dots, k_d m)$. Also, $\mathbf{k} \leq \mathbf{n}$ (resp. $\mathbf{k} \geq \mathbf{n}$) means $k_i \leq n_i$ (resp. $k_i \geq n_i$), $i = 1, 2, \dots, d$. Denote by $S_{\mathbf{n}} = \sum_{\mathbf{k} \leq \mathbf{n}} X_{\mathbf{k}}$ and $\mathbf{1} = (1, \dots, 1) \in \mathbb{N}^d$. It is known that, if $\{X_{\mathbf{n}}\}$ is a field of i.i.d.r.v.s, then

$$\limsup_{\mathbf{n} \rightarrow \infty} \frac{|S_{\mathbf{n}}|}{(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} = (EX_{\mathbf{1}}^2)^{1/2} \quad a.s. \quad (1.3)$$

if and only if $EX_1 = 0$ and $EX_1^2 \log^{d-1}(|X_1|)/\log \log(|X_1|) < \infty$, where $\mathbf{n} \rightarrow \infty$ means $n_1 \rightarrow \infty, \dots, n_d \rightarrow \infty$. When $\{X_{\mathbf{n}}; \mathbf{n} \in \mathbb{N}^d\}$ is a negatively associated field of random variables, Zhang and Wen (2001a) and Zhang and Wang (1999) established the weak convergence, the law of large numbers and the complete convergence similar to those for fields of independent random variables. This paper is to establish the law of the iterated logarithm similar to (1.3) for a negatively associated random field.

Theorem 1.1 *Let $d \geq 2$ be a positive integer, and $\{X_{\mathbf{n}}; \mathbf{n} \in \mathbb{N}^d\}$ be a weakly stationary negatively associated field of identically distributed random variables satisfying*

$$EX_1 = 0 \quad \text{and} \quad EX_1^2 \log^{d-1}(|X_1|)/\log \log(|X_1|) < \infty. \quad (1.4)$$

Denote by $\Upsilon(\mathbf{j} - \mathbf{i}) = \text{Cov}(X_{\mathbf{j}}, X_{\mathbf{i}})$ and $\sigma^2 = \sum_{\mathbf{j} \in \mathbb{Z}^d} \Upsilon(\mathbf{j})$. Then

$$\limsup_{\mathbf{n} \rightarrow \infty} \frac{S_{\mathbf{n}}}{(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} = \sigma \quad a.s. \quad (1.5)$$

Theorem 1.2 *Let $d \geq 2$ be a positive integer, and $\{X_{\mathbf{n}}; \mathbf{n} \in \mathbb{N}^d\}$ be a negatively associated field of identically distributed random variables satisfying (1.4). Then*

$$\limsup_{\mathbf{n} \rightarrow \infty} \frac{|S_{\mathbf{n}}|}{(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} \leq (EX_1^2)^{1/2} \quad a.s.$$

The following theorem tells us that the condition (1.4) is necessary for the law of the iterated logarithm to hold.

Theorem 1.3 *Let $d \geq 1$ be a positive integer, and $\{X_{\mathbf{n}}; \mathbf{n} \in \mathbb{N}^d\}$ be a negatively associated field of identically distributed random variables. If*

$$P \left(\limsup_{\mathbf{n} \rightarrow \infty} \frac{|S_{\mathbf{n}}|}{(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} < \infty \right) > 0, \quad (1.6)$$

then (1.4) holds.

In showing the law of the iterated logarithm, a main step is to establish the exponential inequalities. In the case of $d = 1$, such exponential inequalities for negatively associated random variables are established by Shao (2000) by using a comparison theorem between negatively associated random variables and independent random variables. However, if $d \geq 2$, such a comparison theorem fails for the maximum partial sums (cf., Bulinski and

Suquet 2001). In section 2, we establish a Kolmogorov type exponential inequality as well as a moment inequality of the maximum partial sums of a negatively associated field via a different method. Theorems 1.1-1.3 are proved in Section 3.

2 Moment inequalities and exponential inequalities

First, we have the following moment inequalities and exponential inequalities for the partial sums.

Lemma 2.1 *Let $p \geq 2$ and let $\{Y_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$ be a negatively associated field of random variables with $EY_{\mathbf{k}} = 0$ and $E|Y_{\mathbf{k}}|^p < \infty$. Then*

$$E\left|\sum_{i \leq \mathbf{n}} Y_i\right|^p \leq 2(15p/\ln p)^p \left\{ \left(\sum_{i \leq \mathbf{n}} EY_i^2 \right)^{p/2} + \sum_{i \leq \mathbf{n}} E|Y_i|^p \right\}.$$

Lemma 2.2 *Let $\{Y_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$ be a negatively associated field of random variables with zero means and finite second moments. Let $T_{\mathbf{k}} = \sum_{i \leq \mathbf{k}} Y_i$ and $B_{\mathbf{n}}^2 = \sum_{\mathbf{k} \leq \mathbf{n}} EY_{\mathbf{k}}^2$. Then for all $x > 0$ and $a > 0$,*

$$P(T_{\mathbf{n}} \geq x) \leq P\left(\max_{\mathbf{k} \leq \mathbf{n}} Y_{\mathbf{k}} > a\right) + \exp\left(-\frac{x^2}{2(ax + B_{\mathbf{n}}^2)}\right). \quad (2.1)$$

Lemma 2.3 *Let $\{Y_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$ be a negatively associated field of random variables with $EY_{\mathbf{k}} = 0$ and $E|Y_{\mathbf{k}}|^3 < \infty$. Denote by $T_{\mathbf{k}} = \sum_{i \leq \mathbf{k}} Y_i$ and $B_{\mathbf{n}}^2 = \sum_{\mathbf{k} \leq \mathbf{n}} EY_{\mathbf{k}}^2$. Then for all $x > 0$,*

$$\begin{aligned} P(T_{\mathbf{n}} \geq xB_{\mathbf{n}}) &\geq (1 - \Phi(x+1)) - 6B_{\mathbf{n}}^{-2} \sum_{1 \leq i \neq j \leq \mathbf{n}} |E(Y_i Y_j)| \\ &\quad - 12B_{\mathbf{n}}^{-3} \sum_{\mathbf{k} \leq \mathbf{n}} E|Y_{\mathbf{k}}|^3 \end{aligned} \quad (2.2)$$

where $\Phi(x)$ is the distribution of a standard normal variable.

Proofs of Lemmas 2.1- 2.3: In the case of $d = 1$, Lemma 2.1 is proved by Shao (2000), and Lemma 2.3 is proved by Shao and Su (1999). Also, (2.1) follows from the following inequality easily:

$$P(T_{\mathbf{n}} \geq x) \leq P\left(\max_{\mathbf{k} \leq \mathbf{n}} Y_{\mathbf{k}} > a\right) + \exp\left\{\frac{x}{a} - \left(\frac{x}{a} + \frac{B_{\mathbf{n}}^2}{a^2}\right) \ln\left(\frac{xa}{B_{\mathbf{n}}^2} + 1\right)\right\}.$$

The later is proved by Su, et al.(1997). Since Lemmas 2.1-2.3 do not involve the partial order of the index set, so they are valid for $d \geq 2$ also. In fact, when $d \geq 2$, there is a

one-one map $\pi : \{\mathbf{k} : \mathbf{k} \leq \mathbf{n}\} \rightarrow \{1, 2, \dots, |\mathbf{n}|\}$. By noting that $\{Y_{\pi^{-1}(i)} : i = 1, \dots, |\mathbf{n}|\}$ is a negatively associated sequence and $\sum_{i=1}^{|\mathbf{n}|}(\cdot) = \sum_{\mathbf{k} \leq \mathbf{n}}(\cdot)$, $\max_{\mathbf{k} \leq \mathbf{n}} Y_{\mathbf{k}} = \max_{i \leq |\mathbf{n}|} Y_{\pi^{-1}(i)}$, the results follow.

In a same way, one can extend (1.2) of shao (2000) to the case of $d \geq 2$.

Lemma 2.4 *Let $\{Y_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$ be a negatively associated field and $\{Y_{\mathbf{k}}^*; \mathbf{k} \leq \mathbf{n}\}$ be a field of independent random variables such that for each $\mathbf{k}$, $Y_{\mathbf{k}}$ and $Y_{\mathbf{k}}^*$ have the same distribution. Then*

$$Ef\left(\sum_{\mathbf{k} \leq \mathbf{n}} Y_{\mathbf{k}}\right) \leq Ef\left(\sum_{\mathbf{k} \leq \mathbf{n}} Y_{\mathbf{k}}^*\right)$$

for any convex function f on $\mathbb{R}$, whenever the expectations exist.

It shall be mentioned that it is impossible to find a one-one map $\pi : \{\mathbf{k} : \mathbf{k} \leq \mathbf{n}\} \rightarrow \{1, 2, \dots, |\mathbf{n}|\}$ such that $\sum_{i=1}^{|\mathbf{m}|} Y_{\pi^{-1}(i)} = \sum_{\mathbf{k} \leq \mathbf{m}} Y_{\mathbf{k}}$ for all $\mathbf{m} \leq \mathbf{n}$. So, the inequalities for maximum partial sums can not be extended directly.

For maximum partial sums, Zhang and Wen (2001a) established two moment inequalities.

Proposition 2.1 (Zhang and Wen 2001a) *Let $p \geq 2$, and let $\{Y_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$ be a negatively associated field of random variables with $EY_{\mathbf{k}} = 0$ and $E|Y_{\mathbf{k}}|^p < \infty$. Suppose that $\{\epsilon_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$ is a field of i.i.d.r.v.s with $P(\epsilon_{\mathbf{k}} = \pm 1) = 1/2$. Also, suppose that $\{\epsilon_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$ is independent of $\{Y_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$. Denote by $T_{\mathbf{k}} = \sum_{i \leq \mathbf{k}} Y_i$, $M_{\mathbf{n}} = \max_{\mathbf{k} \leq \mathbf{n}} |T_{\mathbf{k}}|$, $\tilde{T}_{\mathbf{k}} = \sum_{i \leq \mathbf{k}} Y_i$, $\tilde{M}_{\mathbf{n}} = \max_{\mathbf{k} \leq \mathbf{n}} |\tilde{T}_{\mathbf{k}}|$ and $\|\mathbf{X}\|_p = (E|X|^p)^{1/p}$. Then*

$$\|M_{\mathbf{n}}\|_p \leq 5\|\tilde{M}_{\mathbf{n}}\|_p + \|M_{\mathbf{n}}\|_1, \quad (2.3)$$

and there exists a constant $A_{p,d}$ depending on p and d such that

$$E|M_{\mathbf{n}}|^p \leq A_{p,d} \left\{ (E|M_{\mathbf{n}}|)^p + \left(\sum_{\mathbf{k} \leq \mathbf{n}} EY_{\mathbf{k}}^2 \right)^{p/2} + \sum_{\mathbf{k} \leq \mathbf{n}} E|Y_{\mathbf{k}}|^p \right\}. \quad (2.4)$$

Proposition 2.2 (Zhang and Wen 2001a) *Let $\{Y_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$ be a strictly stationary negatively associated field of random variables with $EY_1 = 0$ and $0 < EY_1^2 < \infty$. Denote by $T_{\mathbf{k}} = \sum_{i \leq \mathbf{k}} Y_i$. Then there exists a constant K , depending only on d , such that*

$$\limsup_{n \rightarrow \infty} |\mathbf{n}|^{-1} E \max_{\mathbf{k} \leq \mathbf{n}} T_{\mathbf{k}}^2 \leq K EY_1^2. \quad (2.5)$$

The next theorem gives the estimate of EM_n in (2.4) for a non-stationary negatively associated field.

Theorem 2.1 *Let $\{Y_k; k \leq n\}$ be a negatively associated field of random variables with $EY_k = 0$ and $EY_k^2 < \infty$. Denote by $T_k = \sum_{i \leq k} Y_i$, $M_n = \max_{k \leq n} |T_k|$. Then there is a constant C_d depending only on d such that*

$$EM_n \leq C_d \sqrt{\sum_{k \leq n} EY_k^2}. \quad (2.6)$$

Proof. We will prove (2.6) by induction on d . We use the argument of Utev and Peligrad (2003). For each n , define

$$a_n = \sup_Y \left(E \max_{m \leq n} \left| \sum_{k \leq m} Y_k \right| / \left[\sum_{k \leq n} EY_k^2 \right]^{1/2} \right) \quad (2.7)$$

where the supremum is taken over all fields $Y := \{Y_i\}$ of square integrable centered negatively associated random variables.

Fix such a random field $\{Y_i\}$ and in addition without loss of generality assume that

$$\sum_{k \leq n} EY_k^2 = 1.$$

Let M be a positive integer that will be specified later. Let $f(x) = ((-M^{-1/2}) \vee x) \wedge M^{-1/2}$. For $k \leq n$ define: $\xi_k = f(Y_k) - Ef(Y_k)$ and $\eta_k = Y_k - \xi_k$. Then both $\{\xi_n\}$ and $\{\eta_n\}$ are negatively associated fields. Since

$$\sum_{k \leq n} E|\eta_k| \leq 2M^{1/2} \sum_{k \leq n} EY_k^2 = 2M^{1/2},$$

we get

$$E \max_{m \leq n} \left| \sum_{k \leq m} Y_k \right| \leq E \max_{m \leq n} \left| \sum_{k \leq m} \xi_k \right| + 2M^{1/2}.$$

To estimate $E \max_{m \leq n} \left| \sum_{k \leq m} \xi_k \right|$, we shall use a blocking procedure.

Write $\vec{i} = (i_1, \dots, i_{d-1})$. Take $m_0 = 0$ and define the integers m_k recursively by

$$m_k = \min \left\{ m : m > m_{k-1}, \sum_{j=m_{k-1}+1}^m \sum_{\vec{i} \leq \vec{n}} E\xi_{i,j}^2 > \frac{1}{M} \right\}.$$

Note that, if we denote by ℓ the number of integers produced by this procedure, i.e.: $m_0, m_1, \dots, m_{\ell-1}$, we have

$$1 \geq \sum_{k=1}^{\ell-1} \sum_{j=m_{k-1}+1}^{m_k} \sum_{\bar{i}, j} \mathbb{E} \xi_{\bar{i}, j}^2 > \frac{\ell-1}{M} \quad \text{so that } \ell \leq M.$$

Write $S_{\bar{m}, k} = \sum_{j=m_{k-1}+1}^{m_k} \sum_{\bar{i} \leq \bar{m}} \xi_{\bar{i}, j}$ for $1 \leq k \leq \ell-1$ and for convenience, $m_\ell = n_d$ that is $S_{\bar{m}, \ell} = \sum_{j=m_{\ell-1}+1}^{n_d} \sum_{\bar{i} \leq \bar{m}} \xi_{\bar{i}, j}$. It is obvious that

$$\begin{aligned} & \mathbb{E} \max_{\bar{m} \leq \bar{n}} \left| \sum_{\mathbf{k} \leq \bar{m}} \xi_{\mathbf{k}} \right| \\ & \leq \mathbb{E} \max_{1 \leq k \leq \ell} \max_{\bar{m} \leq \bar{n}} \left| \sum_{j=1}^k S_{\bar{m}, j} \right| + \mathbb{E} \max_{1 \leq k \leq \ell} \max_{\bar{m} \leq \bar{n}} \left(\max_{m_{k-1} < j < m_k} \left| \sum_{t=m_{k-1}+1}^j \sum_{\bar{\mathbf{k}} \leq \bar{m}} \xi_{\bar{\mathbf{k}}, t} \right| \right) \\ & =: I + II. \end{aligned} \tag{2.8}$$

We evaluate the two terms in the right hand side of (2.8) separately.

By the induction hypothesis and Cauchy-Schwartz inequality

$$\begin{aligned} I & \leq \sum_{k=1}^{\ell} \mathbb{E} \max_{\bar{m} \leq \bar{n}} |S_{\bar{m}, k}| \leq C_{d-1} \sum_{k=1}^{\ell} \sqrt{\sum_{\bar{i} \leq \bar{n}} \mathbb{E} \left(\sum_{j=m_{k-1}+1}^{m_k} \xi_{\bar{i}, j} \right)^2} \\ & \leq C_{d-1} \sum_{k=1}^{\ell} \sqrt{\sum_{j=m_{k-1}+1}^{m_k} \sum_{\bar{i} \leq \bar{n}} \mathbb{E} \xi_{\bar{i}, j}^2} \leq C_{d-1} M^{1/2} \sqrt{\sum_{\bar{i} \leq \bar{n}} \mathbb{E} \xi_{\bar{i}}^2} \leq K_{d-1} M^{1/2}. \end{aligned}$$

When $d = 1$, the above inequality obviously holds with $C_0 = 1$, since $\max_{\bar{m} \leq \bar{n}}$ does not appear. To estimate II we notice that

$$(II)^4 \leq \sum_{k=1}^{\ell} \mathbb{E} \max_{\bar{m} \leq \bar{n}} \max_{m_{k-1} < j < m_k} \left| \sum_{t=m_{k-1}+1}^j \sum_{\bar{\mathbf{k}} \leq \bar{m}} \xi_{\bar{\mathbf{k}}, t} \right|^4.$$

By (2.4), we obtain

$$\begin{aligned} & \mathbb{E} \max_{\bar{m} \leq \bar{n}} \max_{m_{k-1} < j < m_k} \left| \sum_{t=m_{k-1}+1}^j \sum_{\bar{\mathbf{k}} \leq \bar{m}} \xi_{\bar{\mathbf{k}}, t} \right|^4 \\ & \leq A_{4,d} \left\{ \mathbb{E}^4 \max_{\bar{m} \leq \bar{n}} \max_{m_{k-1} < j < m_k} \left| \sum_{t=m_{k-1}+1}^j \sum_{\bar{\mathbf{k}} \leq \bar{m}} \xi_{\bar{\mathbf{k}}, t} \right|^4 \right. \\ & \quad \left. + \sum_{t=m_{k-1}+1}^{m_k-1} \sum_{\bar{\mathbf{k}} \leq \bar{n}} \mathbb{E} \xi_{\bar{\mathbf{k}}, t}^4 + \left(\sum_{t=m_{k-1}+1}^{m_k-1} \sum_{\bar{\mathbf{k}} \leq \bar{n}} \mathbb{E} \xi_{\bar{\mathbf{k}}, t}^2 \right)^2 \right\}. \end{aligned}$$

By the definition of m_k ,

$$\sum_{t=m_{k-1}+1}^{m_k-1} \sum_{\bar{\mathbf{k}} \leq \bar{n}} \mathbb{E} \xi_{\bar{\mathbf{k}}, t}^2 \leq \frac{1}{M},$$

so that by using the notation (2.7) for $a_{\mathbf{n}}$ and the definition of $\xi_{\mathbf{k}}$ we obtain

$$\sum_{t=m_{k-1}+1}^{m_k-1} \sum_{\bar{\mathbf{k}} \leq \bar{\mathbf{n}}} \mathbb{E} \xi_{\bar{\mathbf{k}},t}^2 \leq \frac{4}{M} \sum_{t=m_{k-1}+1}^{m_k-1} \sum_{\bar{\mathbf{k}} \leq \bar{\mathbf{n}}} \mathbb{E} \xi_{\bar{\mathbf{k}},t}^4 \leq \frac{4}{M^2}$$

and

$$\mathbb{E}^4 \max_{\bar{\mathbf{m}} \leq \bar{\mathbf{n}}} \max_{m_{k-1} < j < m_k} \left| \sum_{t=m_{k-1}+1}^j \sum_{\bar{\mathbf{k}} \leq \bar{\mathbf{m}}} \xi_{\bar{\mathbf{k}},t} \right| \leq a_{\mathbf{n}}^4 \left(\sum_{t=m_{k-1}+1}^{m_k-1} \sum_{\bar{\mathbf{k}} \leq \bar{\mathbf{n}}} \mathbb{E} \xi_{\bar{\mathbf{k}},t}^2 \right)^2 \leq \frac{a_{\mathbf{n}}^4}{M^2}.$$

Hence we obtain

$$(II)^4 \leq A_{4,d} \left\{ \frac{a_{\mathbf{n}}^4}{M} + \frac{4}{M} + \frac{1}{M} \right\}.$$

Now by (2.8) and the estimates for I and II we get

$$\mathbb{E} \max_{\bar{\mathbf{m}} \leq \bar{\mathbf{n}}} \left| \sum_{\mathbf{k} \leq \bar{\mathbf{m}}} Y_{\mathbf{k}} \right| \leq (A_{4,d}/M)^{1/4} a_{\mathbf{n}} + (5A_{4,d}/M)^{1/4} + C_{d-1} M^{1/2} + 2M^{1/2}.$$

Therefore, by the definition of $a_{\mathbf{n}}$,

$$a_{\mathbf{n}} \leq (A_{4,d}/M)^{1/4} a_{\mathbf{n}} + (5A_{4,d}/M)^{1/4} + C_{d-1} M^{1/2} + 2M^{1/2}.$$

Letting $M = [16A_{4,d}] + 1$ yields

$$a_{\mathbf{n}} \leq 2 + 2(C_{d-1} + 2)M^{1/2}.$$

The proof is now completed.

Combining (2.4) and (2.6) we obtain the following result on the moment inequality for the maximum partial sums.

Theorem 2.2 *Let $p \geq 2$, and let $\{Y_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$ be a negatively associated field of random variables with $\mathbb{E} Y_{\mathbf{k}} = 0$ and $\mathbb{E} |Y_{\mathbf{k}}|^p < \infty$. Then there exists a constant $C_{p,d}$ depending on p and d such that*

$$\mathbb{E} \max_{\bar{\mathbf{m}} \leq \bar{\mathbf{n}}} \left| \sum_{\mathbf{k} \leq \bar{\mathbf{m}}} Y_{\mathbf{k}} \right|^p \leq C_{p,d} \left\{ \left(\sum_{\mathbf{k} \leq \bar{\mathbf{n}}} \mathbb{E} Y_{\mathbf{k}}^2 \right)^{p/2} + \sum_{\mathbf{k} \leq \bar{\mathbf{n}}} \mathbb{E} |Y_{\mathbf{k}}|^p \right\}. \quad (2.9)$$

In the case of $d = 1$, inequality (2.9) is first obtained by Shao (2000). Before that, Su et al (1997) proved that

$$\mathbb{E} \max_{m \leq n} \left| \sum_{k=1}^m Y_k \right|^p \leq C_p \left\{ \left(n \max_{k \leq n} \mathbb{E} Y_k^2 \right)^{p/2} + n \sup_{k \leq n} \mathbb{E} |Y_k|^p \right\}.$$

With the same proof, one can show that (2.9) of a weakly dependent random field with $\lim_{n \rightarrow \infty} \rho_n^* < 1$ (for definition, see Peligrad and Gut (1999)) and a ρ^- -mixing random field (for definition, see Zhang and Wang (1999) or Zhang and Wen (2001a)).

Now, we begin to establish the following Kolmogorov type exponential inequality for maximum partial sums.

Theorem 2.3 *Let $\{Y_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$ be a negatively associated field of random variables with $EY_{\mathbf{k}} = 0$ and $|Y_{\mathbf{k}}| \leq b$ a.s for some $0 < b < \infty$. Denote by $T_{\mathbf{k}} = \sum_{\mathbf{i} \leq \mathbf{k}} Y_{\mathbf{i}}$, $M_{\mathbf{n}} = \max_{\mathbf{k} \leq \mathbf{n}} |T_{\mathbf{k}}|$ and $B_{\mathbf{n}}^2 = \sum_{\mathbf{k} \leq \mathbf{n}} EY_{\mathbf{k}}^2$. Then for all $x > 0$,*

$$P(M_{\mathbf{n}} - 2EM_{\mathbf{n}} \geq 20x) \leq 2^{d+1} \exp\left(-\frac{x^2}{2(bx + B_{\mathbf{n}}^2)}\right). \quad (2.10)$$

Proof. Let $\{\epsilon_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$ is a field of i.i.d.r.v.s with $P(\epsilon_{\mathbf{k}} = \pm 1) = 1/2$. Also, assume that $\{\epsilon_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$ is independent of $\{Y_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$. Denote by $\tilde{T}_{\mathbf{n}} = \sum_{\mathbf{k} \leq \mathbf{n}} \epsilon_{\mathbf{k}} Y_{\mathbf{k}}$, $T_{\mathbf{n},1} = \sum_{\mathbf{k} \leq \mathbf{n}, \epsilon_{\mathbf{k}}=1} Y_{\mathbf{k}}$ and $T_{\mathbf{n},2} = \sum_{\mathbf{k} \leq \mathbf{n}, \epsilon_{\mathbf{k}}=-1} Y_{\mathbf{k}}$. First, we show that

$$Ee^{M_{\mathbf{n}}} \leq 2^d e^{2\|M_{\mathbf{n}}\|_1} Ee^{10|\tilde{T}_{\mathbf{n}}|} \leq 2^{d+1} e^{2\|M_{\mathbf{n}}\|_1} \prod_{\mathbf{k} \leq \mathbf{n}} Ee^{20\epsilon_{\mathbf{k}} Y_{\mathbf{k}}}. \quad (2.11)$$

By the Lévy inequality, we have

$$P\{\tilde{M}_{\mathbf{n}} \geq x\} = E_Y P_{\epsilon}\{\tilde{M}_{\mathbf{n}} \geq x\} \leq 2^d E_Y P_{\epsilon}\{|\tilde{T}_{\mathbf{n}}| \geq x\} \leq 2^d P\{|\tilde{T}_{\mathbf{n}}| \geq x\}, \quad \forall x \geq 0,$$

where $E_Y(\cdot) = E[\cdot | \epsilon_{\mathbf{k}}, \mathbf{k} \in \mathbb{N}^d]$, and E_{ϵ} , P_{ϵ} etc are defined similarly. Then

$$Ee^{10\tilde{M}_{\mathbf{n}}} \leq 2^d Ee^{10|\tilde{T}_{\mathbf{n}}|}.$$

So, from (2.3) it follows that

$$\begin{aligned} Ee^{M_{\mathbf{n}}} &= 1 + EM_{\mathbf{n}} + \sum_{q=2}^{\infty} \frac{EM_{\mathbf{n}}^q}{q!} \\ &\leq 1 + \|M_{\mathbf{n}}\|_1 + \sum_{q=2}^{\infty} \frac{(5\|\tilde{M}_{\mathbf{n}}\|_q + \|M_{\mathbf{n}}\|_1)^q}{q!} \\ &\leq 1 + \|M_{\mathbf{n}}\|_1 + \sum_{q=2}^{\infty} \frac{(2\|5\tilde{M}_{\mathbf{n}} + \|M_{\mathbf{n}}\|_1\|_q)^q}{q!} \\ &= 1 + \|M_{\mathbf{n}}\|_1 + \sum_{q=2}^{\infty} \frac{E(10\tilde{M}_{\mathbf{n}} + 2\|M_{\mathbf{n}}\|_1)^q}{q!} \\ &\leq 1 + \sum_{q=1}^{\infty} \frac{E(10\tilde{M}_{\mathbf{n}} + 2\|M_{\mathbf{n}}\|_1)^q}{q!} = E\left\{1 + \sum_{q=1}^{\infty} \frac{(10\tilde{M}_{\mathbf{n}} + 2\|M_{\mathbf{n}}\|_1)^q}{q!}\right\} \\ &= Ee^{10\tilde{M}_{\mathbf{n}} + 2\|M_{\mathbf{n}}\|_1} \leq 2^d e^{2\|M_{\mathbf{n}}\|_1} Ee^{10|\tilde{T}_{\mathbf{n}}|}. \end{aligned}$$

Note that

$$\mathbb{E}e^{10\tilde{T}_n} = \mathbb{E}e^{10T_{n,1}-10T_{n,2}} \leq \frac{1}{2}\mathbb{E}(e^{20T_{n,1}} + e^{-20T_{n,2}}).$$

For fixed $\{\epsilon_{\mathbf{k}}; \mathbf{k} \leq \mathbf{n}\}$, we have by Lemma 2.4 or the definition (1.1),

$$\begin{aligned} \mathbb{E}_Y e^{20T_{n,1}} &\leq \prod_{\epsilon_{\mathbf{k}}=1} \mathbb{E}_Y e^{20Y_{\mathbf{k}}} = \prod_{\epsilon_{\mathbf{k}}=1} \mathbb{E}_Y e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}} \\ &\leq \prod_{\epsilon_{\mathbf{k}}=1} \mathbb{E}_Y e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}} \cdot \prod_{\epsilon_{\mathbf{k}}=-1} \mathbb{E}_Y e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}} = \prod_{\mathbf{k} \leq \mathbf{n}} \mathbb{E}_Y e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}}, \end{aligned}$$

since $\mathbb{E}_Y e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}} \geq e^{\mathbb{E}_Y(20\epsilon_{\mathbf{k}}Y_{\mathbf{k}})} = 1$. It follows that

$$\begin{aligned} \mathbb{E}e^{20T_{n,1}} &= \mathbb{E}_{\epsilon} \mathbb{E}_Y e^{20T_{n,1}} \leq \mathbb{E}_{\epsilon} \left(\prod_{\mathbf{k} \leq \mathbf{n}} \mathbb{E}_Y e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}} \right) \\ &= \prod_{\mathbf{k} \leq \mathbf{n}} \mathbb{E}_{\epsilon} \mathbb{E}_Y e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}} = \prod_{\mathbf{k} \leq \mathbf{n}} \mathbb{E}e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}}. \end{aligned}$$

Similarly,

$$\mathbb{E}e^{-20T_{n,2}} \leq \prod_{\mathbf{k} \leq \mathbf{n}} \mathbb{E}e^{-20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}} = \prod_{\mathbf{k} \leq \mathbf{n}} \mathbb{E}e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}}.$$

It follows that

$$\mathbb{E}e^{10\tilde{T}_n} \leq \prod_{\mathbf{k} \leq \mathbf{n}} \mathbb{E}e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}}.$$

Similarly,

$$\mathbb{E}e^{-10\tilde{T}_n} \leq \prod_{\mathbf{k} \leq \mathbf{n}} \mathbb{E}e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}}.$$

(2.11) is proved. Now, from (2.11) it follows that for any $x > 0$ and $t > 0$,

$$\mathbb{P}(M_{\mathbf{n}} - 2\mathbb{E}M_{\mathbf{n}} \geq 20x) \leq e^{-20tx-2t\|M_{\mathbf{n}}\|_1} \mathbb{E}e^{tM_{\mathbf{n}}} \leq 2^{d+1}e^{-20tx} \prod_{\mathbf{k} \leq \mathbf{n}} \mathbb{E}e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}}.$$

Since

$$\begin{aligned} \mathbb{E}e^{20\epsilon_{\mathbf{k}}Y_{\mathbf{k}}} &= \mathbb{E}\left\{1 + 20t\epsilon_{\mathbf{k}}Y_{\mathbf{k}} + \frac{e^{20t\epsilon_{\mathbf{k}}Y_{\mathbf{k}}} - 1 - 20t\epsilon_{\mathbf{k}}Y_{\mathbf{k}}}{(\epsilon_{\mathbf{k}}Y_{\mathbf{k}})^2}(\epsilon_{\mathbf{k}}Y_{\mathbf{k}})^2\right\} \\ &\leq 1 + (e^{20tb} - 1 - 20tb)b^{-2}\mathbb{E}Y_{\mathbf{k}}^2 \\ &\leq \exp\{(e^{20tb} - 1 - 20tb)b^{-2}\mathbb{E}Y_{\mathbf{k}}^2\}, \end{aligned}$$

it follows that

$$\mathbb{P}(M_{\mathbf{n}} - 2\mathbb{E}M_{\mathbf{n}} \geq 20x) \leq 2^{d+1}e^{-20tx} \exp\{(e^{20tb} - 1 - 20tb)b^{-2}B_{\mathbf{n}}^2\}.$$

Letting $20t = \frac{1}{b} \log(1 + \frac{xb}{B_n^2})$ yields

$$\begin{aligned} P(M_n - 2EM_n \geq 20x) &\leq 2^{d+1} \exp \left\{ \frac{x}{b} - \left(\frac{x}{b} + \frac{B_n^2}{b^2} \right) \log \left(1 + \frac{xb}{B_n^2} \right) \right\} \\ &\leq 2^{d+1} \exp \left\{ - \frac{x^2}{2(bx + B_n^2)} \right\}. \end{aligned}$$

3 Proofs of the laws of iterated logarithm

We need two more lemmas.

Lemma 3.1 *Let $d \geq 2$ and let $\{Y_{\mathbf{k}}; \mathbf{k} \in \mathbb{N}^d\}$ be a negatively associated field of identically distributed random variables with*

$$EY_1 = 0 \text{ and } EY_1^2 \log^{d-1}(|Y_1|) / \log \log(|Y_1|) < \infty.$$

Denote by $T_n = \sum_{\mathbf{k} \leq \mathbf{n}} Y_{\mathbf{k}}$ and $M_n = \max_{\mathbf{k} \leq \mathbf{n}} |T_{\mathbf{k}}|$. Then

$$\limsup_{n \rightarrow \infty} \frac{M_n}{(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} \leq 20(EY_1^2)^{1/2} \quad a.s. \quad (3.1)$$

Proof. Let $0 < \epsilon < 1$ be an arbitrary but fixed number. Let $b_m = \frac{\epsilon}{40}(EY_1^2)^{1/2}(m / \log \log m)^{1/2}$, $f_{\mathbf{k}}(x) = (-b_{|\mathbf{k}|}) \vee (x \wedge b_{|\mathbf{k}|})$, $g_{\mathbf{k}}(x) = x - f_{\mathbf{k}}(x)$. Define

$$\begin{aligned} \bar{Y}_{\mathbf{k}} &= f_{\mathbf{k}}(Y_{\mathbf{k}}) - Ef_{\mathbf{k}}(Y_{\mathbf{k}}), \quad \hat{Y}_{\mathbf{k}} = g_{\mathbf{k}}(Y_{\mathbf{k}}), \\ \bar{T}_n &= \sum_{\mathbf{k} \leq \mathbf{n}} \bar{Y}_{\mathbf{k}}, \quad \bar{M}_n = \max_{\mathbf{k} \leq \mathbf{n}} |\bar{T}_{\mathbf{k}}|, \quad \hat{T}_n = \sum_{\mathbf{k} \leq \mathbf{n}} \hat{Y}_{\mathbf{k}}. \end{aligned}$$

First, we show that

$$\frac{\hat{T}_n - E\hat{T}_n}{(2|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} \rightarrow 0 \quad a.s. \quad \text{as } \mathbf{n} \rightarrow \infty. \quad (3.2)$$

Since

$$\begin{aligned} \frac{\sum_{\mathbf{k} \leq \mathbf{n}} E|\hat{Y}_{\mathbf{k}}|}{(2|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} &\leq \frac{\sum_{\mathbf{k} \leq \mathbf{n}} E|Y_{\mathbf{k}}| I\{|Y_{\mathbf{k}}| \geq b_{|\mathbf{k}|}\}}{(2|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} \\ &\leq C \frac{\sum_{\mathbf{k} \leq \mathbf{n}} (\frac{\log \log |\mathbf{k}|}{|\mathbf{k}|})^{1/2} EY_1^2 I\{|Y_1| \geq b_{|\mathbf{k}|}\}}{(2|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} \\ &\leq C \frac{\sum_{\mathbf{k} \leq \mathbf{n}} |\mathbf{k}|^{-1/2} EY_1^2 I\{|Y_1| \geq b_{|\mathbf{k}|}\}}{|\mathbf{n}|^{1/2}} \\ &= o(1) \frac{\sum_{\mathbf{k} \leq \mathbf{n}} |\mathbf{k}|^{-1/2}}{|\mathbf{n}|^{1/2}} \rightarrow 0 \quad \text{as } |\mathbf{n}| \rightarrow \infty, \end{aligned} \quad (3.3)$$

it is enough to show that

$$\frac{\sum_{\mathbf{k} \leq \mathbf{n}} |\hat{Y}_{\mathbf{k}}|}{(2\|\mathbf{n}\| \log \log \|\mathbf{n}\|)^{1/2}} \rightarrow 0 \quad a.s. \quad (3.4)$$

For $\mathbf{n} = (n_1, \dots, n_d) \in \mathbb{N}^d$, let $I(\mathbf{n}) = \{\mathbf{k} = (k_1, \dots, k_d) : 2^{n_i-1} \leq k_i \leq 2^{n_i} - 1, i = 1, \dots, d\}$.

(3.4) will be true if we have

$$\frac{\sum_{\mathbf{k} \in I(\mathbf{n})} \hat{Y}_{\mathbf{k}}^+}{(2\|\mathbf{n}\| \log \log 2\|\mathbf{n}\|)^{1/2}} \rightarrow 0 \quad a.s. \quad (3.5)$$

and

$$\frac{\sum_{\mathbf{k} \in I(\mathbf{n})} \hat{Y}_{\mathbf{k}}^-}{(2\|\mathbf{n}\| \log \log 2\|\mathbf{n}\|)^{1/2}} \rightarrow 0 \quad a.s. \quad (3.6)$$

We show (3.5) only since (3.6) can be showed similarly. Let $\alpha_m := \alpha(m) = (2m \log \log m)^{1/2}$ and $Z_{\mathbf{k}} = (\hat{Y}_{\mathbf{k}}^+) \wedge \alpha_{|\mathbf{k}|}$. It is easily seen that $Z_{\mathbf{k}} = 0$ if $Y_{\mathbf{k}} \leq b_{|\mathbf{k}|}$, $Z_{\mathbf{k}} = Y_{\mathbf{k}} - b_{|\mathbf{k}|}$ if $b_{|\mathbf{k}|} \leq Y_{\mathbf{k}} \leq b_{|\mathbf{k}|} + \alpha_{|\mathbf{k}|}$ and $Z_{\mathbf{k}} = \alpha_{|\mathbf{k}|}$ if $Y_{\mathbf{k}} \geq b_{|\mathbf{k}|} + \alpha_{|\mathbf{k}|}$. Also, $\{Z_{\mathbf{k}}; \mathbf{k} \in \mathbb{N}^d\}$ is a negatively associated field of random variables. Obviously,

$$\begin{aligned} \sum_{\mathbf{n}} \mathbb{P}(\hat{Y}_{\mathbf{n}}^+ \neq Z_{\mathbf{n}}) &\leq \sum_{\mathbf{n}} \mathbb{P}(Y_{\mathbf{n}} \geq \alpha_{|\mathbf{n}|}) \leq C \sum_{m=1}^{\infty} (\log m)^{d-1} \mathbb{P}(Y_1 \geq \alpha_m) \\ &\leq C \mathbb{E} Y_1^2 \log^{d-1}(|Y_1|) / \log \log(|Y_1|) < \infty. \end{aligned}$$

Also by (3.3), using the notation $2^{\mathbf{n}} = (2^{n_1}, \dots, 2^{n_d})$ for $\mathbf{n} = (n_1, \dots, n_d) \in \mathbb{N}^d$, one has that,

$$\frac{|\sum_{\mathbf{k} \in I(\mathbf{n})} \mathbb{E} Z_{\mathbf{k}}|}{(2\|\mathbf{n}\| \log \log 2\|\mathbf{n}\|)^{1/2}} \leq \frac{\sum_{\mathbf{k} \leq 2^{\mathbf{n}}} \mathbb{E} |\hat{Y}_{\mathbf{k}}|}{(2\|\mathbf{n}\| \log \log 2\|\mathbf{n}\|)^{1/2}} \rightarrow 0.$$

So, (3.5) is equivalent to

$$\frac{\sum_{\mathbf{k} \in I(\mathbf{n})} (Z_{\mathbf{k}} - \mathbb{E} Z_{\mathbf{k}})}{(2\|\mathbf{n}\| \log \log 2\|\mathbf{n}\|)^{1/2}} \rightarrow 0 \quad a.s. \quad (3.7)$$

Let

$$\Lambda(\mathbf{n}) = \sum_{\mathbf{k} \in I(\mathbf{n})} \frac{\mathbb{E} Y_1^2 I\{b_{|\mathbf{k}|} < |Y_1| \leq 2\alpha_{|\mathbf{k}|}\}}{\alpha_{|\mathbf{k}|}^2}.$$

Note that $b_{|\mathbf{k}|} < \alpha_{|\mathbf{k}|}$ if $|\mathbf{k}|$ is large enough. From Lemma 2.1, it follows that for $\|\mathbf{n}\|$ large enough and any $\delta > 0$,

$$\begin{aligned}
& \mathbb{P}\left(\left|\sum_{\mathbf{k} \in I(\mathbf{n})} (Z_{\mathbf{k}} - \mathbb{E}Z_{\mathbf{k}})\right| \geq \delta(2^{\|\mathbf{n}\|} \log \log 2^{\|\mathbf{n}\|})^{1/2}\right) \\
& \leq C \frac{\mathbb{E}\left|\sum_{\mathbf{k} \in I(\mathbf{n})} (Z_{\mathbf{k}} - \mathbb{E}Z_{\mathbf{k}})\right|^4}{(2^{\|\mathbf{n}\|} \log \log 2^{\|\mathbf{n}\|})^2} \\
& \leq C(\alpha(2^{\|\mathbf{n}\|}))^{-4} \left\{ \left(\sum_{\mathbf{k} \in I(\mathbf{n})} \mathbb{E}|Z_{\mathbf{k}}|^2 \right)^2 + \sum_{\mathbf{k} \in I(\mathbf{n})} \mathbb{E}|Z_{\mathbf{k}}|^4 \right\} \\
& \leq C(\alpha(2^{\|\mathbf{n}\|}))^{-4} \left\{ \left(\sum_{\mathbf{k} \in I(\mathbf{n})} \mathbb{E}Y_1^2 I\{b_{|\mathbf{k}|} < |Y_1| \leq \alpha_{|\mathbf{k}|} + b_{|\mathbf{k}|}\} \right)^2 \right. \\
& \quad \left. + \sum_{\mathbf{k} \in I(\mathbf{n})} \mathbb{E}Y_1^4 I\{b_{|\mathbf{k}|} < |Y_1| \leq \alpha_{|\mathbf{k}|} + b_{|\mathbf{k}|}\} \right\} \\
& \quad + C(\alpha(2^{\|\mathbf{n}\|}))^{-4} \left\{ \left(\sum_{\mathbf{k} \in I(\mathbf{n})} \alpha_{|\mathbf{k}|}^2 \mathbb{P}(|Y_1| \geq \alpha_{|\mathbf{k}|} + b_{|\mathbf{k}|}) \right)^2 \right. \\
& \quad \left. + \sum_{\mathbf{k} \in I(\mathbf{n})} \alpha_{|\mathbf{k}|}^4 \mathbb{P}(|Y_1| \geq \alpha_{|\mathbf{k}|} + b_{|\mathbf{k}|}) \right\} \\
& \leq C \left\{ \Lambda^2(\mathbf{n}) + \sum_{\mathbf{k} \in I(\mathbf{n})} \frac{\mathbb{E}Y_1^4 I\{b_{|\mathbf{k}|} < |Y_1| \leq 2\alpha_{|\mathbf{k}|}\}}{\alpha_{|\mathbf{k}|}^4} \right\} \\
& \quad + C \left\{ \left(\sum_{\mathbf{k} \in I(\mathbf{n})} \mathbb{P}(|Y_1| \geq \alpha_{|\mathbf{k}|}) \right)^2 + \sum_{\mathbf{k} \in I(\mathbf{n})} \mathbb{P}(|Y_1| \geq \alpha_{|\mathbf{k}|}) \right\}.
\end{aligned}$$

Obviously,

$$\begin{aligned}
& \sum_{\mathbf{n}} \left\{ \left(\sum_{\mathbf{k} \in I(\mathbf{n})} \mathbb{P}(|Y_1| \geq \alpha_{|\mathbf{k}|}) \right)^2 + \sum_{\mathbf{k} \in I(\mathbf{n})} \mathbb{P}(|Y_1| \geq \alpha_{|\mathbf{k}|}) \right\} \\
& \leq C \sum_{\mathbf{n}} \mathbb{P}(|Y_1| \geq \alpha_{|\mathbf{n}|}) \leq C \mathbb{E}Y_1^2 \log^{d-1}(|Y_1|) / \log \log(|Y_1|) < \infty.
\end{aligned}$$

Also,

$$\begin{aligned}
& \sum_{\mathbf{n}} \sum_{\mathbf{k} \in I(\mathbf{n})} \frac{\mathbb{E}Y_1^4 I\{b_{|\mathbf{k}|} < |Y_1| \leq 2\alpha_{|\mathbf{k}|}\}}{\alpha_{|\mathbf{k}|}^4} \leq \sum_{\mathbf{n}} \frac{\mathbb{E}Y_1^4 I\{b_{|\mathbf{n}|} < |Y_1| \leq 2\alpha_{|\mathbf{n}|}\}}{\alpha_{|\mathbf{n}|}^4} \\
& \leq \sum_{m=1}^{\infty} (\log m)^{d-1} \frac{\mathbb{E}Y_1^4 I\{|Y_1| \leq 2\alpha_m\}}{\alpha_m^4} \\
& \leq C \sum_{m=1}^{\infty} (\log m)^{d-1} \sum_{k=1}^m \frac{\mathbb{E}Y_1^4 I\{2\alpha_{k-1} < |Y_1| \leq 2\alpha_k\}}{\alpha_m^4} \\
& \leq C \sum_{k=1}^{\infty} \sum_{m=k}^{\infty} \frac{(\log m)^{d-1}}{(m \log \log m)^2} \mathbb{E}Y_1^4 I\{2\alpha_{k-1} < |Y_1| \leq 2\alpha_k\} \\
& \leq C \sum_{k=1}^{\infty} \frac{(\log k)^{d-1}}{k(\log \log k)^2} (k \log \log k) \mathbb{E}Y_1^2 I\{2\alpha_{k-1} < |Y_1| \leq 2\alpha_k\} \\
& \leq C \mathbb{E}Y_1^2 \log^{d-1}(|Y_1|) / \log \log(|Y_1|) < \infty,
\end{aligned}$$

and, similarly to (3.8) of Li and Wu (1989) we have

$$\sum_{\mathbf{n}} \Lambda^2(\mathbf{n}) < \infty.$$

It follows that for arbitrary $\delta > 0$,

$$\sum_{\mathbf{n}} \mathbb{P}\left(\left|\sum_{\mathbf{k} \in I(\mathbf{n})} (Z_{\mathbf{k}} - \mathbb{E}Z_{\mathbf{k}})\right| \geq \delta(2^{\|\mathbf{n}\|} \log \log 2^{\|\mathbf{n}\|})^{1/2}\right) < \infty,$$

which implies (3.7) by the Borel-Cantelli lemma. Thus (3.2) holds.

Now, by applying (2.10) to $x = (1 + 2\epsilon)(\mathbb{E}Y_1^2)^{1/2}(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}$ and $b = 2b_{|\mathbf{n}|}$, it follows that

$$\begin{aligned} & \mathbb{P}(\overline{M}_{\mathbf{n}} - 2\mathbb{E}\overline{M}_{\mathbf{n}} \geq 20(1 + 2\epsilon)(\mathbb{E}Y_1^2)^{1/2}(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}) \\ & \leq 2^{d+1} \exp\{-(1 + \epsilon)d \log \log |\mathbf{n}|\} \leq 2^{d+1} (\log |\mathbf{n}|)^{-(1+\epsilon)d}. \end{aligned}$$

For $\theta > 1$ and $\mathbf{m} \in \mathbb{N}^d$, let $\mathbf{N}_{\mathbf{m}} = ([\theta^{m_1}], \dots, [\theta^{m_d}])$. It follows that

$$\begin{aligned} & \sum_{\mathbf{m}} \mathbb{P}(\overline{M}_{\mathbf{N}_{\mathbf{m}}} - 2\mathbb{E}\overline{M}_{\mathbf{N}_{\mathbf{m}}} \geq 20(1 + 2\epsilon)(\mathbb{E}Y_1^2)^{1/2}(2d|\mathbf{N}_{\mathbf{m}}| \log \log |\mathbf{N}_{\mathbf{m}}|)^{1/2}) \\ & \leq C \sum_{\mathbf{m}} \|\mathbf{m}\|^{-(1+\epsilon)d} \leq C \sum_{i=1}^{\infty} i^{d-1} i^{-(1+\epsilon)d} < \infty. \end{aligned}$$

Notice $\mathbb{E}\overline{M}_{\mathbf{N}_{\mathbf{m}}} \leq C|\mathbf{N}_{\mathbf{m}}|^{1/2}$ by Theorem 2.1. From the Borel-Cantelli lemma, it follows that

$$\begin{aligned} & \limsup_{\mathbf{m} \rightarrow \infty} \frac{\overline{M}_{\mathbf{N}_{\mathbf{m}}}}{(2d|\mathbf{N}_{\mathbf{m}}| \log \log |\mathbf{N}_{\mathbf{m}}|)^{1/2}} \\ & \leq 20(1 + 2\epsilon)(\mathbb{E}Y_1^2)^{1/2} + 2 \limsup_{\mathbf{m} \rightarrow \infty} \frac{\mathbb{E}\overline{M}_{\mathbf{N}_{\mathbf{m}}}}{(2d|\mathbf{N}_{\mathbf{m}}| \log \log |\mathbf{N}_{\mathbf{m}}|)^{1/2}} \\ & = 20(1 + 2\epsilon)(\mathbb{E}Y_1^2)^{1/2}. \end{aligned}$$

So, we have

$$\begin{aligned} & \limsup_{\mathbf{n} \rightarrow \infty} \frac{\overline{M}_{\mathbf{n}}}{(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} \\ & \leq \limsup_{\mathbf{n} \rightarrow \infty} \max_{\mathbf{N}_{\mathbf{m}-1} \leq \mathbf{n} \leq \mathbf{N}_{\mathbf{m}}} \frac{\overline{M}_{\mathbf{N}_{\mathbf{m}}}}{(2d|\mathbf{N}_{\mathbf{m}-1}| \log \log |\mathbf{N}_{\mathbf{m}-1}|)^{1/2}} \\ & \leq 20\theta^{d/2}(1 + 2\epsilon)(\mathbb{E}Y_1^2)^{1/2} \quad a.s. \end{aligned} \tag{3.8}$$

Finally, from (3.2) and (3.8) it follows that (3.1) holds.

Lemma 3.2 *Let $\{X_{\mathbf{k}}; \mathbf{k} \in \mathbb{N}^d\}$ be a negatively associated field of bounded random variables with $EX_{\mathbf{k}} = 0$ for all $\mathbf{k} \in \mathbb{N}^d$. Denote by $S_{\mathbf{n}} = \sum_{\mathbf{k} \leq \mathbf{n}} X_{\mathbf{k}}$. Then for any $\delta > 0$,*

$$\limsup_{\mathbf{n} \rightarrow \infty} \frac{\max_{\mathbf{k} \leq \delta \mathbf{n}} |S_{\mathbf{n}+\mathbf{k}} - S_{\mathbf{n}}|}{(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} \leq 80d(\delta(1+2\delta)^d)^{1/2} \sup_{\mathbf{k}} (EX_{\mathbf{k}}^2)^{1/2} \quad (3.9)$$

Proof. Assume that $|X_{\mathbf{k}}| \leq b$ a.s. with $0 < b < \infty$. Denote $\beta = \sup_{\mathbf{k}} (EX_{\mathbf{k}}^2)^{1/2}$. From Theorem 2.2 it follows that

$$\mathbb{E} \max_{\mathbf{m} \leq \mathbf{n}} \left(\sum_{\mathbf{k} \leq \mathbf{m}} X_{\mathbf{k}} \right)^2 \leq C|\mathbf{n}|\beta^2. \quad (3.10)$$

For fixed $\mathbf{n}$ and $\mathbf{m} \leq \mathbf{n}$, let $Y_{\mathbf{k}} = 0$ for $\mathbf{k} \leq \mathbf{m}$ and $Y_{\mathbf{k}} = X_{\mathbf{k}}$ otherwise, and let $T_{\mathbf{k}} = \sum_{\mathbf{i} \leq \mathbf{k}} Y_{\mathbf{i}}$ for $\mathbf{k} \leq \mathbf{m} + \delta \mathbf{n}$, $B_{\mathbf{m}+[\delta \mathbf{n}]}^2 = \sum_{\mathbf{k} \leq \mathbf{m}+[\delta \mathbf{n}]} \mathbb{E} Y_{\mathbf{k}}^2$. Then for $\mathbf{m} \leq \mathbf{n}$,

$$B_{\mathbf{m}+[\delta \mathbf{n}]}^2 \leq (|\mathbf{m} + [\delta \mathbf{n}]| - |\mathbf{m}|)\beta^2 \leq \delta d(1+\delta)^d |\mathbf{n}|\beta^2.$$

So by Theorem 2.3,

$$\begin{aligned} & \mathbb{P} \left(\max_{\mathbf{k} \leq \delta \mathbf{n}} |S_{\mathbf{m}+\mathbf{k}} - S_{\mathbf{m}}| - 2\mathbb{E} \max_{\mathbf{k} \leq \delta \mathbf{n}} |S_{\mathbf{m}+\mathbf{k}} - S_{\mathbf{m}}| \geq 20x \right) \\ &= \mathbb{P} \left(\max_{\mathbf{k} \leq \mathbf{m}+[\delta \mathbf{n}]} |T_{\mathbf{k}}| - \mathbb{E} \max_{\mathbf{k} \leq \mathbf{m}+[\delta \mathbf{n}]} |T_{\mathbf{k}}| \geq 20x \right) \\ &\leq 2^{d+1} \exp \left\{ - \frac{x^2}{2(bx + B_{\mathbf{m}+[\delta \mathbf{n}]}^2)} \right\} \\ &\leq 2^{d+1} \exp \left\{ - \frac{x^2}{2(bx + \delta d(1+\delta)^d |\mathbf{n}|\beta^2)} \right\}. \end{aligned} \quad (3.11)$$

Note that by (3.10) we have

$$\begin{aligned} & \max_{\mathbf{m} \leq \mathbf{n}} \mathbb{E} \max_{\mathbf{k} \leq \delta \mathbf{n}} |S_{\mathbf{m}+\mathbf{k}} - S_{\mathbf{m}}| \leq 2\mathbb{E} \max_{\mathbf{k} \leq \mathbf{n}+[\delta \mathbf{n}]} |S_{\mathbf{k}}| \leq 2(\mathbb{E} \max_{\mathbf{k} \leq \mathbf{n}+[\delta \mathbf{n}]} S_{\mathbf{k}}^2)^{1/2} \\ &\leq K(|\mathbf{n} + [\delta \mathbf{n}]|\beta^2)^{1/2} \leq K(1+\delta)^{d/2} |\mathbf{n}|^{1/2} \beta. \end{aligned}$$

From (3.11) it follows that for $|\mathbf{n}|$ large enough

$$\begin{aligned} & \max_{\mathbf{m} \leq \mathbf{n}} \mathbb{P} \left(\max_{\mathbf{k} \leq \delta \mathbf{n}} |S_{\mathbf{m}+\mathbf{k}} - S_{\mathbf{m}}| \geq 20d(1+2\epsilon)(2\delta(1+\delta)^d d^2 \beta^2 |\mathbf{n}| \log \log |\mathbf{n}|)^{1/2} \right) \\ &\leq 2^{d+1} \exp \left\{ - (1+\epsilon)d \log \log |\mathbf{n}| \right\}. \end{aligned}$$

Now, let $I_{\mathbf{p}} = \{\mathbf{p} + \mathbf{k} : \mathbf{k} \leq [\delta \mathbf{n}]\}$. Then there are at most $[(\delta^{-1} + 1)^d] + 1$ such $I_{\mathbf{p}}$ s whose union covers $\{\mathbf{k} : \mathbf{k} \leq \mathbf{n}\}$. It follows that

$$\begin{aligned}
& \mathbb{P}\left(\max_{\mathbf{m} \leq \mathbf{n}} \max_{\mathbf{k} \leq \delta \mathbf{n}} |S_{\mathbf{m}+\mathbf{k}} - S_{\mathbf{m}}| \geq (1+2\epsilon)80d(\delta(1+2\delta)^d \beta^2 |\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}\right) \\
& \leq \left(\left(\frac{1}{\delta} + 1\right)^d + 1\right) \max_{\mathbf{p}} \mathbb{P}\left(\max_{\mathbf{m} \in I_{\mathbf{p}}} \max_{\mathbf{k} \leq \delta \mathbf{n}} |S_{\mathbf{m}+\mathbf{k}} - S_{\mathbf{m}}| \geq 40d(1+2\epsilon)(4\delta(1+2\delta)^d \beta^2 |\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}\right) \\
& \leq 4\left(\frac{1}{\delta} + 1\right)^d \max_{\mathbf{m} \leq \mathbf{n}} \mathbb{P}\left(\max_{\mathbf{k} \leq 2\delta \mathbf{n}} |S_{\mathbf{m}+\mathbf{k}} - S_{\mathbf{m}}| \geq 20d(1+2\epsilon)(4\delta(1+2\delta)^d \beta^2 |\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}\right) \\
& \leq 4\left(\frac{1}{\delta} + 1\right)^d 2^{d+1} \exp\left(-(1+\epsilon)d \log \log |\mathbf{n}|\right).
\end{aligned}$$

If we choose $\mathbf{n}_{\mathbf{p}} = ([\theta^{p_1}], \dots, [\theta^{p_d}])$, then the sum of the above probability is finite. And then by the Borel-Cantelli lemma,

$$\limsup_{\mathbf{p} \rightarrow \infty} \frac{\max_{\mathbf{m} \leq \mathbf{n}_{\mathbf{p}}} \max_{\mathbf{k} \leq \delta \mathbf{n}_{\mathbf{p}}} |S_{\mathbf{m}+\mathbf{k}} - S_{\mathbf{m}}|}{(2d|\mathbf{n}_{\mathbf{p}}| \log \log |\mathbf{n}_{\mathbf{p}}|)^{1/2}} \leq 80d(\delta(1+2\delta)^d)^{1/2} \beta \quad a.s.,$$

which implies (3.9) easily.

Now, we turn to the

Proof of Theorem 1.1: We can assume $\sigma > 0$, for otherwise, we can consider the field $\{X_{\mathbf{n}} + \epsilon Z_{\mathbf{n}}; \mathbf{n} \in \mathbb{N}^d\}$ instead, where $\{Z_{\mathbf{n}}; \mathbf{n} \in \mathbb{N}^d\}$ is a field of i.i.d. standard normal random variables and $\epsilon > 0$ is an arbitrary number. First we show that

$$\limsup_{\mathbf{n} \rightarrow \infty} \frac{|S_{\mathbf{n}}|}{(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} \leq \sigma \quad a.s. \quad (3.12)$$

For $b > 0$, let $g_b(x) = (-b) \vee (x \wedge b)$ and $h_b(x) = x - g_b(x)$. Then $g_b(x)$ and $h_b(x)$ are both non-decreasing functions of x . Let $\overline{X}_{\mathbf{k}} = g_b(X_{\mathbf{k}}) - \mathbb{E}g_b(X_{\mathbf{k}})$, $\widehat{X}_{\mathbf{k}} = h_b(X_{\mathbf{k}}) - \mathbb{E}h_b(X_{\mathbf{k}})$, $\overline{S}_{\mathbf{n}} = \sum_{\mathbf{k} \leq \mathbf{n}} \overline{X}_{\mathbf{k}}$ and $\overline{M}_{\mathbf{n}} = \max_{\mathbf{k} \leq \mathbf{n}} |\overline{S}_{\mathbf{k}}|$. And define $\widehat{S}_{\mathbf{k}}$, $\widehat{M}_{\mathbf{n}}$ similarly. Then $X_{\mathbf{k}} = \overline{X}_{\mathbf{k}} + \widehat{X}_{\mathbf{k}}$ and, $\{\overline{X}_{\mathbf{k}}; \mathbf{k} \in \mathbb{N}^d\}$ and $\{\widehat{X}_{\mathbf{k}}; \mathbf{k} \in \mathbb{N}^d\}$ are both negatively associated fields of identically distributed random variables with $\mathbb{E}\overline{X}_{\mathbf{k}} = \mathbb{E}\widehat{X}_{\mathbf{k}} = 0$ and $|\overline{X}_{\mathbf{k}}| \leq 2b$. Then by Lemma 3.1,

$$\begin{aligned}
& \limsup_{\mathbf{n} \rightarrow \infty} \frac{|\widehat{S}_{\mathbf{n}}|}{(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} \\
& \leq 20(\mathbb{E}\widehat{X}_1^2)^{1/2} \leq 20(\mathbb{E}X_1^2 I\{|X_1| \geq b\})^{1/2} \rightarrow 0 \quad a.s. \quad \text{as } b \rightarrow \infty.
\end{aligned} \quad (3.13)$$

So it suffices to show that for any $\epsilon > 0$,

$$\limsup_{\mathbf{n} \rightarrow \infty} \frac{|\overline{S}_{\mathbf{n}}|}{(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} \leq (1+\epsilon)^2 \sigma \quad a.s. \quad (3.14)$$

if b is large enough.

Let $I_m = \{\mathbf{k} : 1 \leq k_i \leq m, i = 1, \dots, d\}$ and $Y_{\mathbf{k}} = \sum_{i \in I_m} \bar{X}_{(\mathbf{k}-1)m+i}$. Since

$$\mathbb{E} \left(\sum_{i \in I_m} X_{(\mathbf{k}-1)m+i} \right)^2 = \mathbb{E} S_{1m}^2 / m^d \rightarrow \sigma^2$$

and

$$\mathbb{E} \left(Y_{\mathbf{k}} - \sum_{i \in I_m} X_{(\mathbf{k}-1)m+i} \right)^2 / m^d \leq 2\mathbb{E} X_1^2 I\{|X_1| \geq b\} \rightarrow 0 \quad \text{as } b \rightarrow \infty,$$

we can choose b and m large enough such that

$$\sup_{\mathbf{k}} \mathbb{E} Y_{\mathbf{k}}^2 \leq (1 + \epsilon)^2 m^d \sigma^2.$$

For $\theta > 1$, let $\mathbf{N}_{\mathbf{k}} = [\theta^{\mathbf{k}}] =: ([\theta^{k_1}], \dots, [\theta^{k_d}])$. From (2.1), it follows that

$$\mathbb{P}(|\bar{S}_{\mathbf{N}_{\mathbf{k}}m}| \geq x) = \mathbb{P}(|\sum_{i \leq \mathbf{N}_{\mathbf{k}}} Y_i| \geq x) \leq 2 \exp \left\{ - \frac{x^2}{2(2m^d b x + \sum_{i \leq \mathbf{N}_{\mathbf{k}}} \mathbb{E} Y_i^2)} \right\}.$$

Let $x = (1 + \epsilon)^2 \sigma (2dm^d |\mathbf{N}_{\mathbf{k}}| \log \log |\mathbf{N}_{\mathbf{k}}|)^{1/2}$. It follows that

$$\begin{aligned} & \sum_{\mathbf{k}} \mathbb{P}(|\bar{S}_{\mathbf{N}_{\mathbf{k}}m}| \geq (1 + \epsilon)^2 \sigma (2dm^d |\mathbf{N}_{\mathbf{k}}| \log \log |\mathbf{N}_{\mathbf{k}}|)^{1/2}) \\ & \leq \sum_{\mathbf{k}} \mathbb{P}(|\bar{S}_{\mathbf{N}_{\mathbf{k}}m}| \geq (1 + \epsilon)(2d \sum_{i \leq \mathbf{N}_{\mathbf{k}}} \mathbb{E} Y_i^2 \log \log |\mathbf{N}_{\mathbf{k}}|)^{1/2}) \\ & \leq C \sum_{\mathbf{k}} \exp \left\{ - (1 + \epsilon)d \log \log |\mathbf{N}_{\mathbf{k}}| \right\} \\ & \leq C \sum_{\mathbf{k}} \|\mathbf{k}\|^{-(1+\epsilon)d} \leq C \sum_{i=1}^{\infty} i^{d-1} i^{-(1+\epsilon)d} < \infty. \end{aligned}$$

From the Borel-Cantelli lemma, it follows that

$$\begin{aligned} & \limsup_{\mathbf{k} \rightarrow \infty} \frac{|\bar{S}_{\mathbf{N}_{\mathbf{k}}m}|}{(2d |\mathbf{N}_{\mathbf{k}}m| \log \log |\mathbf{N}_{\mathbf{k}}m|)^{1/2}} \\ & = \limsup_{\mathbf{k} \rightarrow \infty} \frac{|\bar{S}_{\mathbf{N}_{\mathbf{k}}m}|}{(2dm^d |\mathbf{N}_{\mathbf{k}}| \log \log |\mathbf{N}_{\mathbf{k}}|)^{1/2}} \leq (1 + \epsilon)^2 \sigma \quad a.s. \end{aligned}$$

So, from Lemma 3.2 it follows that

$$\begin{aligned} & \limsup_{\mathbf{n} \rightarrow \infty} \frac{|\bar{S}_{\mathbf{n}}|}{(2d |\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} \\ & \leq \limsup_{\mathbf{k} \rightarrow \infty} \max_{\mathbf{N}_{\mathbf{k}}m \leq \mathbf{n} \leq \mathbf{N}_{\mathbf{k}+1}m} \frac{|\bar{S}_{\mathbf{n}}|}{(2d |\mathbf{N}_{\mathbf{k}}m| \log \log |\mathbf{N}_{\mathbf{k}}m|)^{1/2}} \\ & \leq \limsup_{\mathbf{k} \rightarrow \infty} \frac{|\bar{S}_{\mathbf{N}_{\mathbf{k}}m}|}{(2d |\mathbf{N}_{\mathbf{k}}m| \log \log |\mathbf{N}_{\mathbf{k}}m|)^{1/2}} \\ & \quad + \limsup_{\mathbf{k} \rightarrow \infty} \max_{\mathbf{N}_{\mathbf{k}}m \leq \mathbf{n} \leq \mathbf{N}_{\mathbf{k}+1}m} \frac{|\bar{S}_{\mathbf{n}} - \bar{S}_{\mathbf{N}_{\mathbf{k}}m}|}{(2d |\mathbf{N}_{\mathbf{k}}m| \log \log |\mathbf{N}_{\mathbf{k}}m|)^{1/2}} \\ & \leq (1 + \epsilon)\sigma + 80d((\theta - 1)(1 + 2(\theta - 1))^d)^{1/2} (\mathbb{E} \bar{X}^2)^{1/2} \quad a.s. \end{aligned}$$

Letting $\theta \rightarrow 1$ completes the proof of (3.14).

Next we show that

$$\limsup_{n \rightarrow \infty} \frac{S_n}{(2d|n| \log \log |n|)^{1/2}} \geq \sigma \quad a.s.$$

Due to (3.13), it suffices to show that for any $0 < \epsilon < 1/9$,

$$\limsup_{n \rightarrow \infty} \frac{\bar{S}_n}{(2d|n| \log \log |n|)^{1/2}} \geq (1 - 9\epsilon)\sigma \quad a.s. \quad (3.15)$$

for b large enough.

For $k \in \mathbb{N}$, let $m_k = \lfloor 2^{k^{1+\epsilon}} \rfloor$, $p_k = \lfloor k^{-2} 2^{k^{1+\epsilon}} \rfloor$, $N_k = (m_k + p_k)k^4$. For $\mathbf{k} \in \mathbb{N}^d$ and an integer q , let $\mathbf{m}_{\mathbf{k}} = (m_{k_1}, \dots, m_{k_d})$, $\mathbf{p}_{\mathbf{k}} = (p_{k_1}, \dots, p_{k_d})$, $\mathbf{N}_{\mathbf{k}} = (N_{k_1}, \dots, N_{k_d})$ and $\mathbf{k}^q = (k_1^q, \dots, k_d^q)$. We first show the following equality

$$\sum_{\mathbf{k}} \mathbb{P}(\bar{S}_{\mathbf{N}_{\mathbf{k}}} \geq (1 - 7\epsilon)\sigma(2d|\mathbf{N}_{\mathbf{k}}| \log \log |\mathbf{N}_{\mathbf{k}}|)^{1/2}) = \infty. \quad (3.16)$$

Set $I_i = I_i(\mathbf{k}) = \{(i-1)(\mathbf{m}_{\mathbf{k}} + \mathbf{p}_{\mathbf{k}}) + \mathbf{1} \leq \mathbf{m} \leq (i-1)(\mathbf{m}_{\mathbf{k}} + \mathbf{p}_{\mathbf{k}}) + \mathbf{m}_{\mathbf{k}}\}$, $A_{\mathbf{k}} = \bigcup_{1 \leq i \leq k^4} I_i$ and $B_{\mathbf{k}} = \{\mathbf{m} : \mathbf{m} \leq \mathbf{N}_{\mathbf{k}}\} \setminus A_{\mathbf{k}}$. Then $\text{Card} A_{\mathbf{k}} = |\mathbf{k}|^4 |\mathbf{m}_{\mathbf{k}}| \sim |\mathbf{N}_{\mathbf{k}}|$ and $\text{Card} B_{\mathbf{k}} = |\mathbf{N}_{\mathbf{k}}| - \text{Card} A_{\mathbf{k}} = o(|\mathbf{N}_{\mathbf{k}}|)$. Let

$$v_{i,1} = \sum_{j \in I_i} \bar{X}_j$$

and

$$S_{\mathbf{k},1} = \sum_{1 \leq i \leq k^4} v_{i,1} = \sum_{j \in A_{\mathbf{k}}} \bar{X}_j, \quad S_{\mathbf{k},2} = \sum_{j \in B_{\mathbf{k}}} \bar{X}_j.$$

Clearly,

$$\bar{S}_{\mathbf{N}_{\mathbf{k}}} = S_{\mathbf{k},1} + S_{\mathbf{k},2}$$

and

$$\sum_{j \in B_{\mathbf{k}}} \mathbb{E} \bar{X}_j^2 / |\mathbf{N}_{\mathbf{k}}| \leq \mathbb{E} X_1^2 \text{Card} B_{\mathbf{k}} / |\mathbf{N}_{\mathbf{k}}| \rightarrow 0, \quad \mathbf{k} \rightarrow \infty.$$

From (2.1), it follows that for any $\epsilon > 0$

$$\begin{aligned} & \sum_{\mathbf{k}} \mathbb{P}(|S_{\mathbf{k},2}| \geq \epsilon \sigma(2d|\mathbf{N}_{\mathbf{k}}| \log \log |\mathbf{N}_{\mathbf{k}}|)^{1/2}) \\ & \leq 2 \sum_{\mathbf{k}} \exp \left\{ - \frac{\epsilon^2 \sigma^2 2d|\mathbf{N}_{\mathbf{k}}| \log \log |\mathbf{N}_{\mathbf{k}}|}{2\{b\epsilon \sigma(2d|\mathbf{N}_{\mathbf{k}}| \log \log |\mathbf{N}_{\mathbf{k}}|)^{1/2} + \sum_{j \in B_{\mathbf{k}}} \mathbb{E} \bar{X}_j^2\}} \right\} \\ & \leq C \sum_{\mathbf{k}} \exp \{ - 3d \log \log |\mathbf{N}_{\mathbf{k}}| \} < \infty. \end{aligned}$$

Thus in order to prove (3.16) is enough to show that

$$\sum_{\mathbf{k}} \mathbb{P}(S_{\mathbf{k},1} \geq (1 - 6\epsilon)\sigma(2d|\mathbf{N}_{\mathbf{k}}| \log \log |\mathbf{N}_{\mathbf{k}}|)^{1/2}) = \infty. \quad (3.17)$$

Let $B_{\mathbf{k}}^2 = \sum_{1 \leq i \leq \mathbf{k}^4} \mathbb{E} v_{i,1}^2$. Since

$$\sum_{1 \leq i \leq \mathbf{k}^4} \mathbb{E} \left(\sum_{j \in I_i} X_j \right)^2 = |\mathbf{k}|^4 \mathbb{E} S_{\mathbf{m}_{\mathbf{k}}}^2 \sim |\mathbf{k}|^4 |\mathbf{m}_{\mathbf{k}}| \sigma^2 \sim |\mathbf{N}_{\mathbf{k}}| \sigma^2$$

and

$$\sum_{1 \leq i \leq \mathbf{k}^4} \mathbb{E} (v_{i,1} - \sum_{j \in I_i} X_j)^2 / |\mathbf{N}_{\mathbf{k}}| \leq C \mathbb{E} X^2 I\{|X| \geq b\} \rightarrow 0 \text{ as } b \rightarrow \infty,$$

for b and $\mathbf{k}$ large enough

$$B_{\mathbf{k}}^2 \geq (1 - \epsilon)^2 |\mathbf{N}_{\mathbf{k}}| \sigma^2.$$

From Lemma 2.3, it follows that

$$\begin{aligned} & \mathbb{P}(S_{\mathbf{k},1} \geq (1 - 6\epsilon)\sigma(2d|\mathbf{N}_{\mathbf{k}}| \log \log |\mathbf{N}_{\mathbf{k}}|)^{1/2}) \\ & \geq \mathbb{P}(S_{\mathbf{k},1} \geq (1 - 5\epsilon)(2dB_{\mathbf{k}}^2|\mathbf{N}_{\mathbf{k}}| \log \log |\mathbf{N}_{\mathbf{k}}|)^{1/2}) \\ & \geq \{1 - \Phi(1 + (1 - 5\epsilon)(2d \log \log |\mathbf{N}_{\mathbf{k}}|)^{1/2})\} - J_{\mathbf{k},1} - J_{\mathbf{k},2} \end{aligned}$$

where

$$\begin{aligned} J_{\mathbf{k},1} &= O(1)|\mathbf{N}_{\mathbf{k}}|^{-1} \sum_{1 \leq i \neq j \leq \mathbf{k}^4} |\mathbb{E}(v_{i,1} v_{j,1})|, \\ J_{\mathbf{k},2} &= O(1)|\mathbf{N}_{\mathbf{k}}|^{-3/2} \sum_{1 \leq i \leq \mathbf{k}^4} \mathbb{E}|v_{i,1}|^3. \end{aligned}$$

Obviously,

$$\begin{aligned} & \sum_{\mathbf{k}} \{1 - \Phi(1 + (1 - 5\epsilon)(2d \log \log |\mathbf{N}_{\mathbf{k}}|)^{1/2})\} \\ & \geq C \sum_{\mathbf{k}} (\log |\mathbf{N}_{\mathbf{k}}|)^{-(1-4\epsilon)d} \geq C \sum_{\mathbf{k}} \|\mathbf{k}\|^{-(1-2\epsilon)d} = \infty, \end{aligned}$$

and by Lemma 2.1,

$$\sum_{\mathbf{k}} J_{\mathbf{k},2} \leq C_p \sum_{\mathbf{k}} |\mathbf{N}_{\mathbf{k}}|^{-3/2} |\mathbf{k}|^4 (|\mathbf{m}_{\mathbf{k}}|^{3/2} (2b)^3 + |\mathbf{m}_{\mathbf{k}}| (2b)^3) \leq C \sum_{\mathbf{k}} |\mathbf{k}|^{-2} < \infty.$$

Also,

$$\begin{aligned} J_{\mathbf{k},1} &= O(1)|\mathbf{N}_{\mathbf{k}}|^{-1} \sum_{1 \leq i \neq j \leq \mathbf{N}_{\mathbf{k}}, j-i \geq p_{\mathbf{k}}} |\text{Cov}(\bar{X}_i, \bar{X}_j)| \\ &= O(1)|\mathbf{N}_{\mathbf{k}}|^{-1} \sum_{1 \leq i \neq j \leq \mathbf{N}_{\mathbf{k}}, j-i \geq p_{\mathbf{k}}} |\text{Cov}(X_i, X_j)| \\ &= O(1) \sum_{p_{\mathbf{k}} \leq j \leq \mathbf{N}_{\mathbf{k}}} |\text{Cov}(X_1, X_{j+1})| = O(1) \sum_{\mathbf{N}_{\mathbf{k}-1} < j \leq \mathbf{N}_{\mathbf{k}}} |\text{Cov}(X_1, X_{j+1})|, \end{aligned}$$

since $p_{k_i} = o(N_{k_i})$ for $i = 1, \dots, d$. It follows that

$$\sum_{\mathbf{k}} J_{\mathbf{k},1} = O(1) \sum_j |\text{Cov}(X_1, X_{j+1})| < \infty.$$

Hence (3.17) is proved.

Now, let $C_{\mathbf{k}} = \{\mathbf{m} : N_{\mathbf{k}-1} < \mathbf{m} \leq N_{\mathbf{k}}\}$, $D_{\mathbf{k}} = \{\mathbf{m} : \mathbf{m} \leq N_{\mathbf{k}}\} \setminus C_{\mathbf{k}}$ and

$$U_{\mathbf{k}} = \sum_{j \in C_{\mathbf{k}}} \overline{X}_j, \quad V_{\mathbf{k}} = \sum_{j \in D_{\mathbf{k}}} \overline{X}_j.$$

Then $\overline{S}_{N_{\mathbf{k}}} = U_{\mathbf{k}} + V_{\mathbf{k}}$, $\text{Card}C_{\mathbf{k}} \sim |N_{\mathbf{k}}|$ and $\text{Card}D_{\mathbf{k}} = o(|N_{\mathbf{k}}|)$. From (2.1), it follows that for any $\delta > 0$,

$$\sum_{\mathbf{k}} \mathbb{P}(|V_{\mathbf{k}}| \geq \delta \sigma(2d|N_{\mathbf{k}}| \log \log |N_{\mathbf{k}}|)^{1/2}) < \infty. \quad (3.18)$$

By the Borel-Cantelli lemma, we conclude that

$$\limsup_{\mathbf{k} \rightarrow \infty} \frac{|V_{\mathbf{k}}|}{(2d|N_{\mathbf{k}}| \log \log |N_{\mathbf{k}}|)^{1/2}} = 0 \quad a.s. \quad (3.19)$$

From (3.16) and (3.18), it follows that

$$\sum_{\mathbf{k}} \mathbb{P}(U_{\mathbf{k}} \geq (1 - 8\epsilon)\sigma(2d|N_{\mathbf{k}}| \log \log |N_{\mathbf{k}}|)^{1/2}) = \infty. \quad (3.20)$$

Note that $\{U_{\mathbf{k}}; \mathbf{k} \in \mathbb{N}^d\}$ is a negatively associated field. It follows that for any x, y and $i \neq j$,

$$\mathbb{P}(U_i \geq x, U_j \geq y) \leq \mathbb{P}(U_i \geq x)\mathbb{P}(U_j \geq y).$$

Hence, by the generalized Borel-Cantelli lemma, (3.20) yields

$$\limsup_{\mathbf{k} \rightarrow \infty} \frac{U_{\mathbf{k}}}{(2d|N_{\mathbf{k}}| \log \log |N_{\mathbf{k}}|)^{1/2}} \geq (1 - 8\epsilon)\sigma \quad a.s. \quad (3.21)$$

From (3.19) and (3.21), it follows that (3.15) holds.

Proof of Theorem 1.2 is similar to that of (3.12) in which we choose $m = 1$ and $Y_{\mathbf{k}} = \overline{X}_{\mathbf{k}}$ instead.

Proof of Theorem 1.3: From (1.6), it follows that there exists a constant $0 < C < \infty$ such that

$$\mathbb{P}\left(\limsup_{n \rightarrow \infty} \frac{|S_{\mathbf{n}}|}{(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} < C\right) > 0.$$

Then

$$\mathbb{P} \left(\limsup_{\mathbf{n} \rightarrow \infty} \frac{|X_{\mathbf{n}}|}{(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}} < 2C \right) > 0. \quad (3.22)$$

Let

$$A_{\mathbf{n}} = \{|X_{\mathbf{n}}| \geq 2C(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}\},$$

$$A_{\mathbf{n}}^+ = \{X_{\mathbf{n}}^+ \geq 2C(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}\},$$

$$A_{\mathbf{n}}^- = \{X_{\mathbf{n}}^- \geq 2C(2d|\mathbf{n}| \log \log |\mathbf{n}|)^{1/2}\}.$$

Note that $\{I\{A_{\mathbf{n}}^+\}; \mathbf{n} \in \mathbb{N}^d\}$ and $\{I\{A_{\mathbf{n}}^-\}; \mathbf{n} \in \mathbb{N}^d\}$ are both negatively associated fields. It follows that for any $\mathbf{m}$ and $\mathbf{n}$,

$$\begin{aligned} \text{Var} \left\{ \sum_{\mathbf{m} \leq \mathbf{k} \leq \mathbf{m} + \mathbf{n}} I\{A_{\mathbf{n}}^+\} \right\} &\leq \sum_{\mathbf{m} \leq \mathbf{k} \leq \mathbf{m} + \mathbf{n}} \mathbb{P}(A_{\mathbf{n}}^+), \\ \text{Var} \left\{ \sum_{\mathbf{m} \leq \mathbf{k} \leq \mathbf{m} + \mathbf{n}} I\{A_{\mathbf{n}}^-\} \right\} &\leq \sum_{\mathbf{m} \leq \mathbf{k} \leq \mathbf{m} + \mathbf{n}} \mathbb{P}(A_{\mathbf{n}}^-). \end{aligned}$$

It follows that

$$\text{Var} \left\{ \sum_{\mathbf{m} \leq \mathbf{k} \leq \mathbf{m} + \mathbf{n}} I\{A_{\mathbf{k}}\} \right\} \leq 2 \sum_{\mathbf{m} \leq \mathbf{k} \leq \mathbf{m} + \mathbf{n}} \mathbb{P}(A_{\mathbf{k}}).$$

Hence from Lemma A.6 of Zhang and Wen (2001b), it follows that for any $\mathbf{m}$

$$(1 - \mathbb{P}(\bigcup_{\mathbf{k} \geq \mathbf{m}} A_{\mathbf{k}}))^2 \sum_{\mathbf{k} \geq \mathbf{m}} \mathbb{P}(A_{\mathbf{k}}) \leq 2\mathbb{P}(\bigcup_{\mathbf{k} \geq \mathbf{m}} A_{\mathbf{k}}). \quad (3.23)$$

Since (3.22) implies $\mathbb{P}(A_{\mathbf{n}}, i.o.) < 1$, we conclude that for some $\mathbf{m}$,

$$\mathbb{P}(\bigcup_{\mathbf{k} \geq \mathbf{m}} A_{\mathbf{k}}) := \beta < 1.$$

Then from (3.23), it follows that

$$\sum_{\mathbf{k} \geq \mathbf{m}} \mathbb{P}(A_{\mathbf{k}}) \leq \frac{2\beta}{(1 - \beta^2)} < \infty.$$

So,

$$\sum_{\mathbf{k} \geq \mathbf{m}} \mathbb{P}(|X_{\mathbf{k}}| \geq 2C(2d|\mathbf{k}| \log \log |\mathbf{k}|)^{1/2}) < \infty,$$

which implies

$$\mathbb{E} X_{\mathbf{1}}^2 \log^{d-1}(|X_{\mathbf{1}}|) / \log \log(|X_{\mathbf{1}}|) < \infty. \quad (3.24)$$

Finally, from (3.24) and the law of the large numbers (c.f. Zhang and Wang 1999), it follows that

$$\lim_n \frac{S_n - |n|EX_1}{|n|} \rightarrow 0 \quad a.s.$$

which together with (1.6) yields $EX_1 = 0$. And then (1.4) holds.

References

- [1] Alam, K. and Saxena, K. M. L. (1981), Positive dependence in multivariate distributions, *Comm. Statist.* **A 10**: 1183-1196.
- [2] Bulinski, A. and Suquet, C. (2001), Normal approximation for quasi-associated random fields, *Statist. Probab. Lett.*, **54**: 215-226.
- [3] Joag-Dev, K. and Proschan, F. (1983), Negative association of random variables, with applications, *Ann Statist.* **11**: 286-295.
- [4] Li, D. L and Wu, Z. Q. (1989), The law of the iterated logarithm for B -Valued random variables with multidimensional indices, *Ann. Probab.* **17**: 760-774.
- [5] Matula, P. (1992), A note on the almost sure convergence of sums of negatively dependent random variables. *Statist. Probab. Lett.* **15**: 209-213.
- [6] Móricz, F. (1983), A general moment inequality for the maximum of rectangular partial sums of multiple series, *Acta. Math. Hungar.* **41**, 337-346.
- [7] Newman, C. M. (1984), Asymptotic independence and limit theorems for positively and negatively dependent random variables. In: Tong, Y. L. (Ed.), *Inequalities in Statistics and Probability*. IMS, Hayward, CA., pp. 127-140.
- [8] Peligrad, M. and Gut, A. (1999), Almost sure results for a class of weak dependent random variables, *J. Theor. Probab.*, **12**: 87-104.

- [9] Roussas, C. G. (1996), Exponential probability inequalities with some applications, In Ferguson, T. S., Shapley, L. S. and MacQueen, J. B. (eds.), *Statistics, Probability and Game Theory*, IMS, Hayward, CA., pp. 303-319.
- [10] Shao, Q. M. (2000), A comparison theorem on maximum inequalities between negatively associated and independent random variables, *J. Theor. Probab.*, **13**: 343-356.
- [11] Shao, Q. M. and Su, C. (1999), The law of the iterated logarithm for negatively associated random variables, *Stochastic Processes & their Appl.* **83**: 139-148.
- [12] Su, C., Zhao, L. C. and Wang, Y. B. (1997), The moment inequalities and weak convergence for negatively associated sequences, *Science in China* **40A**: 172-182.
- [13] Utev, S. and Peligrad, M. (2003), Maximal inequalities and an invariance principle for a class of weakly dependent random variables, *J. Theor. Probab.*, **16**: 101-115.
- [14] Zhang, L. X. and Wang, X. Y. (1999), Convergence rates in the strong laws of asymptotically negatively associated random fields, *Appl. Math.-JCU, Ser. B*, **14**(4): 406-416.
- [15] Zhang, L. X. and Wen J. W. (2001b), A weak convergence for negatively associated fields, *Statist. Probab. Lett.*, **53**:259-267.
- [16] Zhang, L. X. and Wen, J. W. (2001a), Strong laws for sums of B -valued mixing random fields, *Chinese Ann. Math.*, **22A**:205-216. (in Chinese)

LI-XIN ZHANG

DEPARTMENT OF MATHEMATICS

ZHEJIANG UNIVERSITY, YUQUAN CAMPUS

HANGZHOU 310027, ZHEJIANG

P.R. CHINA

EMAIL: STAZLX@ZJU.EDU.HZ.CN